

Research Article

On Finite Interquark Potential in $D = 3$ Driven by a Minimal Length

Patricio Gaete

Departamento de Física and Centro Científico-Tecnológico de Valparaíso, Universidad Técnica Federico Santa María, Valparaíso, Chile

Correspondence should be addressed to Patricio Gaete; patricio.gaete@usm.cl

Received 8 April 2014; Accepted 30 June 2014; Published 17 July 2014

Academic Editor: Stephen Pate

Copyright © 2014 Patricio Gaete. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

We address the effect of a quantum gravity induced minimal length on a physical observable for three-dimensional Yang-Mills. Our calculation is done within stationary perturbation theory. Interestingly enough, we find an ultraviolet finite interaction energy, which contains a regularized logarithmic function and a linear confining potential. This result highlights the role played by the new quantum of length in our discussion.

1. Introduction

It is known that one of the main unsolved problems in high energy physics is a quantitative description (from first principles) of confinement in quantum chromodynamics (QCD). Albeit phenomenological models still represent a key tool for understanding confinement physics. In this context we recall the phenomenon of condensation, where in the scenario of dual superconductivity, it is conjectured that the QCD vacuum behaves as a dual-type II superconductor. More explicitly, due to the condensation of magnetic monopoles, the chromoelectric field acting between $q\bar{q}$ pair is squeezed into strings, and the nonvanishing string tension represents the proportionality constant in the linear potential. Incidentally, lattice calculations have confirmed this picture by showing the formation of tubes of gluonic fields connecting colored charges [1].

It is also known that considerable attention has been paid to the investigation of extensions of the standard model (SM), such as Lorentz invariance violation and fundamental length [2–7], because the SM does not include a quantum theory of gravitation. In this respect we recall that, in the last few years, the emphasis of quantum gravity has been on effective models incorporating a minimal length scale. In fact, there are several approaches on how to incorporate a minimal length scale in a quantum field theory, leading to a model of quantum

space-time [8–11]. Of these, noncommutative quantum field theories have motivated a great interest [12–17]. Notice that this noncommutative geometry is an intrinsic property of space-time. In addition, we also recall that most of the known results in the noncommutative approach have been achieved using a Moyal star-product. Nevertheless, in recent times, a new formulation of noncommutative quantum field theory in the presence of a minimal length has been proposed in [18–20]. Afterwards, this approach was further developed by the introduction of a new multiplication rule, which is known as Voros star-product. Notwithstanding, physics turns out to be independent from the choice of the type of product [21]. Consequently, with the introduction of noncommutativity by means of a minimal length, the theory becomes ultraviolet finite and the cutoff is provided by the noncommutative parameter θ .

In this perspective the present work is an extension of our previous study [22]. Thus, the basic ideas underlying the analysis of this paper are derived from our earlier paper [22]. Specifically, in this work we will focus attention on the impact of a minimal length on a physical observable for pure Yang-Mills theory in $(2 + 1)D$. It is worth noting here that Yang-Mills theories in $(2 + 1)D$ are very relevant for a reliable comparison between results coming from continuum and lattice calculations [23]. Also, $(2 + 1)D$ theories have been raising a great deal of interest in connection with branes activity,

for example, issues like self-duality [24] and new possibilities for supersymmetry breaking as induced by 3-branes [25]. Yet, $(2 + 1)D$ theories may be adopted to describe the high-temperature limit of models in $(3 + 1)D$ [26]. In fact, such theories are of interest to probe low-dimensional condensed matter systems, such as spin or pairing fluctuations by means of effective gauge theories, for which $(2 + 1)D$ theories are a very good approximation [27]. Thus, in order to accomplish the purpose of studying the impact of a minimal length for Yang-Mills theory in $(2 + 1)D$, we will work out the static potential for the case under consideration. As we will see, the presence of a minimal length leads to an ultraviolet finite static potential, which contains a regularized logarithmic function and a linear confining potential. Accordingly, our study offers a straightforward calculation in which some features of three-dimensional nonabelian gauge theories become more transparent.

2. Interaction Energy

We will now discuss the interaction energy for Yang-Mills theory in the presence of a minimal length. We start then with the three-dimensional space-time Lagrangian:

$$\mathcal{L} = -\frac{1}{4} \text{Tr} (F_{\mu\nu} F^{\mu\nu}) = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu}. \quad (1)$$

Here $A_\mu^a(x) = A_\mu^a(x)T^a$ and $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$, with f^{abc} the structure constants of the gauge group.

As we have indicated in [22], our analysis is based in perturbation theory along the lines of [28–30]. To do this, we will work out the vacuum expectation value of the energy operator $H(\langle 0|H|0\rangle)$ at lowest order in g , in the Coulomb gauge. The canonical Hamiltonian can be worked as usual and is given by

$$H = \frac{1}{2} \int d^2x \{ (\mathbf{E}_T^a)^2 + (\mathbf{B}^a)^2 - \phi^a \nabla^2 \phi^a \}, \quad (2)$$

where the color-electric field \mathbf{E}^a has been separated into transverse and longitudinal parts: $\mathbf{E}^a = \mathbf{E}_T^a - \nabla \phi^a$.

Next, by making use of Gauss's law

$$\nabla^2 \phi^a = g (\rho^a - f^{abc} \mathbf{A}^b \cdot \mathbf{E}^c), \quad (3)$$

we get

$$\begin{aligned} \nabla^2 \phi^a &= \left(g \delta^{ap} + g^2 f^{abp} \mathbf{A}^b \cdot \nabla \frac{1}{\nabla^2} \right. \\ &\quad \left. + g^3 f^{abc} \mathbf{A}^b \cdot \nabla \frac{1}{\nabla^2} f^{chp} \mathbf{A}^h \cdot \nabla \frac{1}{\nabla^2} \right) \\ &\quad \times (\rho^p - f^{pde} \mathbf{A}^d \cdot \mathbf{E}_T^e). \end{aligned} \quad (4)$$

Following our earlier procedure, the corresponding formulation of this theory in the presence of a minimal length is by means of a smeared source [22, 31, 32]. Thus, we will take the sources as $\rho^a \equiv \rho_1^a + \rho_2^a = \rho_q^a + \rho_{\bar{q}}^a$, where $\rho_q^a(\mathbf{x}) = t_q^a e^{(\theta/2)\nabla^2} \delta^{(3)}(\mathbf{x} - \mathbf{y}')$ and $\rho_{\bar{q}}^a(\mathbf{x}) = t_{\bar{q}}^a e^{(\theta/2)\nabla^2} \delta^{(3)}(\mathbf{x} - \mathbf{y})$,

where $t_{\bar{q}}^a$ and t_q^a are the color charges of a heavy antiquark \bar{q}_i and a quark q_i in a normalized color singlet state $|\Psi\rangle = N^{-1/2} |q_i\rangle |\bar{q}_i\rangle$. Hence $t_q^a t_{\bar{q}}^a = (1/N) \text{tr}(T^a T^a) = -C_F$, where the anti-Hermitian generators T^a are in the fundamental representation of $SU(N)$.

By proceeding in the same way as in [22], we obtain the expectation value of the energy operator H to order g^2 and g^4 :

$$V = V_1 + V_2, \quad (5)$$

where

$$V_1 = -g^2 \int d^2x \langle 0 | \rho_1^a \frac{1}{\nabla^2} \rho_2^a | 0 \rangle, \quad (6)$$

V_2

$$= -3g^4 f^{abc} f^{chq} \int d^2x \langle 0 | \rho_1^a \frac{1}{\nabla^2} \mathbf{A}^b \cdot \nabla \frac{1}{\nabla^2} \mathbf{A}^h \cdot \nabla \frac{1}{\nabla^2} \rho_2^q | 0 \rangle. \quad (7)$$

The V_1 term is exactly the one obtained in [32]. Consequently, (6) takes the form

$$\begin{aligned} V_1 &= -g^2 C_F \int \frac{d^2k}{(2\pi)^2} \frac{e^{-\theta k^2}}{k^2} e^{-i\mathbf{k}\cdot\mathbf{r}} \\ &= \frac{g^2 C_F}{2\pi} \left\{ \ln(\mu r) - e^{-r^2/4\theta} \ln\left(\frac{r}{2\sqrt{\theta}}\right) \right\}, \end{aligned} \quad (8)$$

with $|\mathbf{r}| \equiv |\mathbf{y} - \mathbf{y}'| = r$ and μ is an infrared regulator. Again, as in our previous analysis [32], unexpected features are found. Interestingly, it is observed that, unlike the Coulomb potential which is singular at the origin, V_1 is finite there: $V_1 = (g^2 C_F / 2\pi) \ln(2\mu\sqrt{\theta})$.

We now turn our attention to the V_2 term, which is given by

$$V_2 = 3g^4 C_A C_F \int \frac{d^2k}{(2\pi)^2} \frac{e^{-\theta k^2}}{k^2} e^{-i\mathbf{k}\cdot\mathbf{r}} \mathcal{J}(\mathbf{k}), \quad (9)$$

where

$$\mathcal{J}(\mathbf{k}) = \int \frac{d^2p}{(2\pi)^2} \frac{1}{2|\mathbf{p}|(\mathbf{p} - \mathbf{k})^2} \left(1 - \frac{(\mathbf{p} \cdot \mathbf{k})^2}{\mathbf{p}^2 \mathbf{k}^2} \right). \quad (10)$$

In passing we recall that to obtain (9) we have expressed the A^{ai} -fields in terms of a normal mode expansion: $A^{ai}(\mathbf{x}, t) = \int (d^3p / \sqrt{(2\pi)^3 2\omega_p}) \sum_\lambda \varepsilon^i(\mathbf{p}, \lambda) [a^a(\mathbf{p}, \lambda) e^{-ipx} + a^{\dagger a}(\mathbf{p}, \lambda) e^{ipx}]$, along with $[a^a(\mathbf{p}, \lambda), a^{\dagger b}(\mathbf{l}, \sigma)] = \delta^{ab} \delta^{\lambda\sigma} \delta^{(3)}(\mathbf{p} - \mathbf{l})$ and $\sum_\lambda \varepsilon^i(\mathbf{k}, \lambda) \varepsilon^j(\mathbf{l}, \lambda) = \delta^{ij} - k^i k^j / \mathbf{k}^2$. Here we would mention that the correction term of order g^4 represents an antiscreening effect. Incidentally, it is of interest to notice that precisely this term is in the origin of asymptotic freedom in the $(3+1)D$ case, which is due to the instantaneous Coulomb interaction of the quarks.

When the integral (10) is performed, one gets

$$\mathcal{J}(k) = \frac{1}{8\pi^2} \frac{1}{|\mathbf{k}|} \left(-1.5706 + \frac{\pi}{2} \right). \quad (11)$$

Expression (9) then becomes

$$V_2 = -\frac{3}{8\pi^2} \left(1.5706 - \frac{\pi}{2}\right) g^4 C_A C_F \int \frac{d^2 k}{(2\pi)^2} \frac{e^{-\theta k^2}}{|\mathbf{k}|^3} e^{-i\mathbf{k}\cdot\mathbf{r}}. \quad (12)$$

We now proceed to calculate the integral (12). Following our earlier procedure [32], (12) is further rewritten as

$$\begin{aligned} I &\equiv \lim_{\varepsilon \rightarrow 0} \tilde{I} \\ &= \lim_{\varepsilon \rightarrow 0} (\mu^2)^{-\varepsilon/2} \int \frac{d^{2+\varepsilon} k}{(2\pi)^2} \frac{e^{-\theta k^2}}{|\mathbf{k}|^3} e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= \lim_{\varepsilon \rightarrow 0} \frac{(\mu^2)^{-\varepsilon/2}}{\Gamma(3/2)} \int_0^\infty dx x^{1/2} \int \frac{d^{2+\varepsilon} k}{(2\pi)^2} e^{-(\theta+x)\mathbf{k}^2} e^{i\mathbf{k}\cdot\mathbf{r}}. \end{aligned} \quad (13)$$

Then, the I term takes the form

$$\begin{aligned} I &= \lim_{\varepsilon \rightarrow 0} \frac{(\mu^2)^{-\varepsilon/2} (r^2)^{-\varepsilon/2}}{\Gamma(3/2) (\pi)^{1+\varepsilon/2}} \\ &\quad \times \frac{1}{4} \int_0^{r^2/4\theta} d\tau \tau^{\varepsilon/2-1} e^{-\tau} \left(\frac{r^2}{4\tau} - \theta\right)^{1/2}. \end{aligned} \quad (14)$$

Hence, at leading order in θ , (14) reduces to

$$\begin{aligned} I &= -\frac{1}{2(\pi)^{3/2}} \left\{ r\gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) + 2\sqrt{\theta} e^{-r^2/4\theta} \right. \\ &\quad \left. + \frac{\theta}{r} \gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) \right\}, \end{aligned} \quad (15)$$

where $\gamma(1/2, r^2/4\theta)$ is the lower incomplete Gamma function defined by the following integral representation:

$$\gamma\left(\frac{a}{b}, x\right) \equiv \int_0^x \frac{du}{u} u^{a/b} e^{-u}. \quad (16)$$

Accordingly, the V_2 term reads

$$\begin{aligned} V_2 &= \frac{3}{16\pi^{7/2}} \left(1.5706 - \frac{\pi}{2}\right) g^4 C_A C_F \\ &\quad \times \left\{ r\gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) + 2\sqrt{\theta} e^{-r^2/4\theta} + \frac{\theta}{r} \gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) \right\}. \end{aligned} \quad (17)$$

Now we focus on the (g^4) screening contribution to the potential, which is due to the exchange of transverse gluons. From our above perturbation theory, we find that V_2^* is given by

$$\begin{aligned} V_2^* &= 2g^4 f_{abc} f_{def} \sum_{n=2} \frac{1}{E_n} \int d^3 x \int d^3 w \langle 0 | \rho_2^a \frac{1}{\sqrt{2}} \mathbf{A}^b \cdot \mathbf{E}_T^c | n \rangle_x \\ &\quad \times \langle n | \rho_1^d \frac{1}{\sqrt{2}} \mathbf{A}^e \cdot \mathbf{E}_T^f | 0 \rangle_w. \end{aligned} \quad (18)$$

In passing we recall that $\langle 0 | \rho_2^a (1/\nabla^2) \mathbf{A}^b \cdot \mathbf{E}_T^c | n \rangle$ is the matrix element in the basis of states in which the nonperturbated Hamiltonian is diagonal. Next, it should be noted that the intermediate state $|n\rangle$ must contain a pair of transverse gluons, since the terms $\mathbf{A}^b \cdot \mathbf{E}_T^c$ must create and destroy dynamical gluon pairs. We can, therefore, write two gluon states as

$$\begin{aligned} &\sum_{n=2} |n\rangle \langle n| \\ &= \frac{1}{2} \sum_{kl} \sum_{\lambda\sigma} \int d^3 k \int d^3 l a^{\dagger e}(\mathbf{k}, \lambda) a^{\dagger f}(\mathbf{l}, \sigma) |0\rangle \langle 0| \\ &\quad \times a^f(\mathbf{l}, \sigma) a^e(\mathbf{k}, \lambda). \end{aligned} \quad (19)$$

By substituting (19) into (18) and following our earlier procedure, the V_2^* term assumes the form

$$V_2^* = -C_A C_F g^4 \int \frac{d^2 k}{(2\pi)^2} \frac{e^{-\theta k^2}}{\mathbf{k}^4} e^{i\mathbf{k}\cdot\mathbf{r}} \mathcal{F}(\mathbf{k}), \quad (20)$$

where

$$\begin{aligned} \mathcal{F}(\mathbf{k}) &= \int \frac{d^2 l}{(2\pi)^2} \frac{(|\mathbf{l}| - |\mathbf{l} - \mathbf{k}|)^2}{4 |\mathbf{l}| |\mathbf{l} - \mathbf{k}| (|\mathbf{l}| + |\mathbf{l} - \mathbf{k}|)} \\ &\quad \times \left(1 - \frac{\mathbf{k}^2}{(\mathbf{l} - \mathbf{k})^2} + \frac{(\mathbf{k} \cdot \mathbf{l})^2}{(\mathbf{l} - \mathbf{k})^2 l^2} \right). \end{aligned} \quad (21)$$

Integrating now over \mathbf{l} , one then obtains $\mathcal{F}(\mathbf{k}) = |\mathbf{k}|((1/2\pi)(15/16) - (1/8) + (3/32\pi^2))$. As a consequence, the V_2^* term becomes

$$V_2^* = -\left(\frac{1}{2\pi} \frac{15}{16} - \frac{1}{8} + \frac{3}{32\pi^2}\right) g^4 C_A C_F \int \frac{d^2 k}{(2\pi)^2} \frac{e^{-\theta k^2}}{|\mathbf{k}|^3} e^{-i\mathbf{k}\cdot\mathbf{r}}. \quad (22)$$

It is straightforward to see that this integral is exactly the one obtained in expression (12).

By putting together (8), (12), and (22), we evaluate the interquark potential in position space. We thus finally obtain

$$\begin{aligned} V(r) &= \frac{g^2 C_F}{2\pi} \left\{ \ln(\mu r) - e^{-r^2/4\theta} \ln\left(\frac{r}{2\sqrt{\theta}}\right) \right\} \\ &\quad + \frac{g^4 C_A C_F}{2(\pi)^{3/2}} 0.165 \\ &\quad \times \left\{ r\gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) + 2\sqrt{\theta} e^{-r^2/4\theta} + \frac{\theta}{r} \gamma\left(\frac{1}{2}, \frac{r^2}{4\theta}\right) \right\}, \end{aligned} \quad (23)$$

which is ultraviolet finite (Figure 1). An immediate consequence of this is that for $\theta = 0$ one obtains the known interquark potential at order g^4 [30]. Note that in Figure 1, for illustrative purposes, we have defined $g^2 C_F / 2\pi = 1$, $g^4 C_A C_F / 2(\pi)^{3/2} = 1$, $\mu = 1$, and $\sqrt{\theta} = 1.4$.




FIGURE 1: The potential V , as a function of $x = r/2\sqrt{\theta}$.

3. Conclusion

To conclude, let us put our work in its proper perspective. As already anticipated, this work is a sequel to [22], where we have considered a three-dimensional extension of the recently $(3 + 1)D$ calculation in the presence of a minimal length. To do this, we have exploited a correct identification of field degrees with observable quantities. Once the identification has been made, the computation of the potential is achieved by means of Gauss' law. Interestingly enough, it was found that the static potential profile is ultraviolet finite, which contains a regularized logarithmic function and a linear potential leading to confinement of static sources. Finally, we note that our results agree for the $\theta = 0$ case with the calculation shown in [30]. Also very recently, in the context of the Georgi-Glashow model it has been shown that there is confinement at distances much larger than the screening length [33]. Since our calculation has shown that there is confinement in three-dimensional Yang-Mills, it seems a challenging work to extend the above analysis to the Georgi-Glashow model. We expect to report on progress along these lines soon.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The author would like to thank José A. Helayël-Neto for useful discussions. This work was partially supported by Fondecyt (Chile) Grant no. 1130426. The author also wishes to thank the Field Theory Group of the CBPF for hospitality and PCI/MCT for support.

References

- [1] S. Capstick et al., "Key issues in hadronic physics," <http://arxiv.org/abs/hep-ph/0012238>.
- [2] G. Amelino-Camelia, "Special treatment," *Nature*, vol. 418, no. 6893, pp. 34–35, 2002.
- [3] T. Jacobson, S. Liberati, and D. Mattingly, "Threshold effects and Planck scale Lorentz violation: combined constraints from high energy astrophysics," *Physical Review D*, vol. 67, Article ID 124011, 2003.
- [4] T. J. Konopka and S. A. Major, "Observational limits on quantum geometry effects," *New Journal of Physics*, vol. 4, article 57, 2002.
- [5] S. Hossenfelder, "Interpretation of quantum field theories with a minimal length scale," *Physical Review D*, vol. 73, Article ID 105013, 2006.
- [6] P. Nicolini, "Noncommutative black holes, the final appeal to quantum gravity: a review," *International Journal of Modern Physics A*, vol. 24, no. 7, p. 1229, 2009.
- [7] P. Nicolini, A. Smailagic, and E. Spallucci, "Noncommutative geometry inspired Schwarzschild black hole," *Physics Letters B*, vol. 632, no. 4, pp. 547–551, 2006.
- [8] J. Magueijo and L. Smolin, "Lorentz invariance with an invariant energy scale," *Physical Review Letters*, vol. 88, no. 19, Article ID 190403, 2002.
- [9] S. Hossenfelder, M. Bleicher, S. Hofmann, J. Ruppert, S. Scherer, and H. Stöcker, "Signatures in the Planck regime," *Physics Letters B: Nuclear, Elementary Particle and High-Energy Physics*, vol. 575, no. 1-2, pp. 85–99, 2003.
- [10] P. Nicolini and E. Spallucci, "Un-spectral dimension and quantum spacetime phases," *Physics Letters B*, vol. 695, no. 1–4, pp. 290–293, 2011.
- [11] M. Sprenger, P. Nicolini, and M. Bleicher, "Physics on the smallest scales: an introduction to minimal length phenomenology," *European Journal of Physics*, vol. 33, no. 4, pp. 853–862, 2012.
- [12] E. Witten, "Noncommutative geometry and string field theory," *Nuclear Physics B*, vol. 268, no. 2, pp. 253–294, 1986.
- [13] N. Seiberg and E. Witten, "String theory and noncommutative geometry," *Journal of High Energy Physics*, vol. 1999, no. 09, article 032, 1999.
- [14] M. R. Douglas and N. A. Nekrasov, "Noncommutative field theory," *Reviews of Modern Physics*, vol. 73, no. 4, pp. 977–1029, 2001.
- [15] R. J. Szabo, "Quantum field theory on noncommutative spaces," *Physics Reports A: Review Section of Physics Letters*, vol. 378, no. 4, pp. 207–299, 2003.
- [16] J. Gomis, K. Kamimura, and T. Mateos, "Gauge and BRST generators for space-time non-commutative U(1) theory," *Journal of High Energy Physics*, vol. 0103, p. 010, 2001.
- [17] A. A. Bichl, J. M. Grimstrup, L. Popp, M. Schweda, and R. Wulkenhaar, "Perturbative analysis of the Seiberg-Witten map," *International Journal of Modern Physics A*, vol. 17, no. 16, pp. 2219–2231, 2002.
- [18] A. Smailagic and E. Spallucci, "UV divergence-free QFT on noncommutative plane," *Journal of Physics A: Mathematical and General*, vol. 36, no. 39, pp. L517–L521, 2003.
- [19] A. Smailagic and E. Spallucci, "Feynman path integral on the non-commutative plane," *Journal of Physics A: Mathematical and General*, vol. 36, no. 33, pp. L467–L471, 2003.
- [20] A. Smailagic and E. Spallucci, "Statistics of equally weighted random paths on a class of self-similar structures," *Journal of Physics A*, vol. 37, no. 1, 2004.
- [21] A. B. Hammou, M. Lagraa, and M. M. Sheikh-Jabbari, "Coherent state induced star product on R_1^3 and the fuzzy sphere," *Physical Review D*, vol. 66, no. 2, Article ID 025025, 11 pages, 2002.

- [22] P. Gaete, “Remarks on the interquark potential in the presence of a minimal length,” *Journal of Physics A*, vol. 46, no. 47, Article ID 475402, 2013.
- [23] C. Feuchter and H. Reinhardt, “Yang-Mills vacuum in Coulomb gauge in $D = 2 + 1$ dimensions,” *Physical Review D: Particles, Fields, Gravitation, and Cosmology*, vol. 77, no. 8, Article ID 085023, 2008.
- [24] H. Singh, “3-branes on Eguchi–Hanson 6D instantons,” *General Relativity and Gravitation*, vol. 39, no. 6, pp. 839–848, 2007.
- [25] O. Bergman, A. Hanany, A. Karch, and B. Kol, “Branes and supersymmetry breaking in 3D gauge theories,” *Journal of High Energy Physics*, vol. 10, article 36, 1999.
- [26] A. Das, *Finite Temperature Field Theory*, World Scientific Publishing, Singapore, 1999.
- [27] D. V. Khveshchenko, “Ghost excitonic insulator transition in layered graphite,” *Physical Review Letters*, vol. 87, Article ID 246802, 2001.
- [28] S. D. Drell, SLAC-PUB-2694, 1981.
- [29] V. N. Gribov, *Lecture at the 12th Winter School of the Leningrad Nuclear Physics Institute*, SLAC-TRANS-176, 1977, Edited by H. D. I. Abarbane.
- [30] E. Bagan, M. Lavelle, and D. McMullan, “The structure of the QCD potential in 2+1 dimensions,” *Physics Letters B*, vol. 477, no. 1–3, pp. 355–360, 2000.
- [31] P. Gaete and E. Spallucci, “Finite axionic electrodynamics from a new non-commutative approach,” *Journal of Physics A*, vol. 45, no. 6, Article ID 065401, 2012.
- [32] P. Gaete, J. Helayel-Neto, and E. Spallucci, “Aspects of finite electrodynamics in $D = 3$ dimensions,” *Journal of Physics A: Mathematical and Theoretical*, vol. 45, no. 21, Article ID 215401, 2012.
- [33] M. M. Anber, “The abelian confinement mechanism revisited: new aspects of the Georgi-GLAshow model,” *Annals of Physics*, vol. 341, pp. 21–55, 2014.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

