

UNIVERSAL SERIES BY TRIGONOMETRIC SYSTEM IN WEIGHTED L^1_μ SPACES

S. A. EPISKOPOSIAN

Received 17 April 2006; Accepted 9 November 2006

We consider the question of existence of trigonometric series universal in weighted $L^1_\mu[0, 2\pi]$ spaces with respect to rearrangements and in usual sense.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

Let X be a Banach space.

Definition 1.1. A series

$$\sum_{k=1}^{\infty} f_k, \quad f_k \in X, \quad (1.1)$$

is said to be universal in X with respect to rearrangements, if for any $f \in X$ the members of (1.1) can be rearranged so that the obtained series $\sum_{k=1}^{\infty} f_{\sigma(k)}$ converges to f by norm of X .

Definition 1.2. The series (1.1) is said to be universal (in X) in the usual sense, if for any $f \in X$ there exists a growing sequence of natural numbers n_k such that the sequence of partial sums with numbers n_k of the series (1.1) converges to f by norm of X .

Definition 1.3. The series (1.1) is said to be universal (in X) concerning partial series, if for any $f \in X$ it is possible to choose a partial series $\sum_{k=1}^{\infty} f_{n_k}$ from (1.1), which converges to the f by norm of X .

Note that many papers are devoted (see [1–10]) to the question on existence of various types of universal series in the sense of convergence *almost everywhere and on a measure*.

The first usual universal in the sense of convergence almost everywhere trigonometric series were constructed by Menshov [6] and Kozlov [5]. The series of the form

$$\frac{1}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx \quad (1.2)$$

2 Universal series by trigonometric system

was constructed just by them such that for any measurable-on- $[0, 2\pi]$ function $f(x)$ there exists the growing sequence of natural numbers n_k such that the series (1.2) having the sequence of partial sums with numbers n_k converges to $f(x)$ almost everywhere on $[0, 2\pi]$. (Note here that in this result, when $f(x) \in L^1_{[0, 2\pi]}$, it is impossible to replace convergence almost everywhere by convergence in the metric $L^1_{[0, 2\pi]}$).

This result was distributed by Talaljan on arbitrary orthonormal complete systems (see [8]). He also established (see [9]) that if $\{\phi_n(x)\}_{n=1}^\infty$ —the normalized basis of space $L^p_{[0, 1]}$, $p > 1$, then there exists a series of the form

$$\sum_{k=1}^{\infty} a_k \phi_k(x), \quad a_k \rightarrow 0, \quad (1.3)$$

which has property: for any measurable function $f(x)$ the members of series (1.3) can be rearranged so that the again received series converge on a measure on $[0, 1]$ to $f(x)$.

Orlicz [7] observed the fact that there exist functional series that are universal with respect to rearrangements in the sense of a.e. convergence in the class of a.e. finite measurable functions.

It is also useful to note that even Riemann proved that every convergent numerical series which is not absolutely convergent is universal with respect to rearrangements in the class of all real numbers.

Let $\mu(x)$ be a measurable-on- $[0, 2\pi]$ function with $0 < \mu(x) \leq 1$, $x \in [0, 2\pi]$, and let $L^1_\mu[0, 2\pi]$ be a space of measurable functions $f(x)$, $x \in [0, 2\pi]$, with

$$\int_0^{2\pi} |f(x)| \mu(x) dx < \infty. \quad (1.4)$$

Grigorian constructed a series of the form (see [3])

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} \quad \text{with} \quad \sum_{k=-\infty}^{\infty} |C_k|^q < \infty \quad \forall q > 2, \quad (1.5)$$

which is universal in $L^1_\mu[0, 2\pi]$ concerning partial series for some weighted function $\mu(x)$, $0 < \mu(x) \leq 1$, $x \in [0, 2\pi]$.

In [2] it is proved that for any given sequence of natural numbers $\{\lambda_m\}_{m=1}^\infty$ with $\lambda_m \nearrow^\infty$ there exists a series by trigonometric system of the form

$$\sum_{k=1}^{\infty} C_k e^{ikx}, \quad C_{-k} = \overline{C}_k, \quad (1.6)$$

with

$$\left| \sum_{k=1}^m C_k e^{ikx} \right| \leq \lambda_m, \quad x \in [0, 2\pi], \quad m = 1, 2, \dots, \quad (1.7)$$

so that for each $\varepsilon > 0$ a weighted function $\mu(x)$,

$$0 < \mu(x) \leq 1, \quad |\{x \in [0, 2\pi] : \mu(x) \neq 1\}| < \varepsilon, \quad (1.8)$$

can be constructed so that the series (1.6) is universal in the weighted space $L_\mu^1[0, 2\pi]$ with respect simultaneously to rearrangements as well as to subseries.

In this paper, we prove the following results.

THEOREM 1.4. *There exists a series of the form*

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} \quad \text{with} \quad \sum_{k=-\infty}^{\infty} |C_k|^q < \infty \quad \forall q > 2 \quad (1.9)$$

such that for any number $\varepsilon > 0$ a weighted function $\mu(x)$, $0 < \mu(x) \leq 1$, with

$$|\{x \in [0, 2\pi] : \mu(x) \neq 1\}| < \varepsilon \quad (1.10)$$

can be constructed so that the series (1.9) is universal in $L_\mu^1[0, 2\pi]$ with respect to rearrangements.

THEOREM 1.5. *There exists a series of the form (1.9) such that for any number $\varepsilon > 0$ a weighted function $\mu(x)$ with (1.10) can be constructed so that the series (1.9) is universal in $L_\mu^1[0, 2\pi]$ in the usual sense.*

2. Basic lemma

LEMMA 2.1. *For any given numbers $0 < \varepsilon < 1/2$, $N_0 > 2$, and a step function*

$$f(x) = \sum_{s=1}^q \gamma_s \cdot \chi_{\Delta_s}(x), \quad (2.1)$$

where Δ_s is an interval of the form $\Delta_m^{(i)} = [(i-1)/2^m, i/2^m]$, $1 \leq i \leq 2^m$, and

$$|\gamma_s| \cdot \sqrt{|\Delta_s|} < \varepsilon^3 \cdot \left(8 \cdot \int_0^{2\pi} f^2(x) dx\right)^{-1}, \quad s = 1, 2, \dots, q, \quad (2.2)$$

there exists a measurable set $E \subset [0, 2\pi]$ and a polynomial $P(x)$ of the form

$$P(x) = \sum_{N_0 \leq |k| < N} C_k e^{ikx}, \quad (2.3)$$

which satisfy the conditions

$$|E| > 2\pi - \varepsilon, \quad (2.4a)$$

$$\int_E |P(x) - f(x)| dx < \varepsilon, \quad (2.4b)$$

$$\sum_{N_0 \leq |k| < N} |C_k|^{2+\varepsilon} < \varepsilon, \quad C_{-k} = \overline{C_k}, \quad (2.4c)$$

$$\max_{N_0 \leq m < N} \left[\int_e \left| \sum_{N_0 \leq |k| \leq m} C_k e^{ikx} \right| dx \right] < \varepsilon + \int_e |f(x)| dx \quad (2.4d)$$

for every measurable subset e of E .

4 Universal series by trigonometric system

Proof. Let $0 < \epsilon < 1/2$ be an arbitrary number.

Set

$$g(x) = 1 \quad \text{if } x \in [0, 2\pi] \setminus \left[\frac{\epsilon \cdot \pi}{2}, \frac{3\epsilon \cdot \pi}{2} \right], \quad (2.5)$$

$$g(x) = 1 - \frac{2}{\epsilon} \quad \text{if } x \in \left[\frac{\epsilon \cdot \pi}{2}, \frac{3\epsilon \cdot \pi}{2} \right]. \quad (2.6)$$

We choose natural numbers ν_1 and N_1 so large that the following inequalities be satisfied:

$$\frac{1}{2\pi} \left| \int_0^{2\pi} g_1(t) e^{-ikt} dt \right| < \frac{\epsilon}{16 \cdot \sqrt{N_0}}, \quad |k| < N_0, \quad (2.7)$$

where

$$g_1(x) = \gamma_1 \cdot g(\nu_1 \cdot x) \cdot \chi_{\Delta_1}(x). \quad (2.8)$$

(By $\chi_E(x)$ we denote the characteristic function of the set E .) We put

$$E_1 = \{x \in \Delta_s : g_s(x) = \gamma_s\}. \quad (2.9)$$

By (2.5), (2.8), and (2.9) we have

$$|E_1| > 2\pi \cdot (1 - \epsilon) \cdot |\Delta_1|, \quad g_1(x) = 0, \quad x \notin \Delta_1, \quad (2.10)$$

$$\int_0^{2\pi} g_1^2(x) dx < \frac{2}{\epsilon} \cdot |\gamma_1|^2 \cdot |\Delta_1|. \quad (2.11)$$

Since the trigonometric system $\{e^{ikx}\}_{k=-\infty}^{\infty}$ is complete in $L^2[0, 2\pi]$, we can choose a natural number $N_1 > N_0$ so large that

$$\int_0^{2\pi} \left| \sum_{0 \leq |k| < N_1} C_k^{(1)} e^{ikx} - g_1(x) \right| dx \leq \frac{\epsilon}{8}, \quad (2.12)$$

where

$$C_k^{(1)} = \frac{1}{2\pi} \int_0^{2\pi} g_1(t) e^{-ikt} dt. \quad (2.13)$$

Hence by (2.7), (2.8), and (2.12) we obtain

$$\int_0^{2\pi} \left| \sum_{N_0 \leq |k| < N_1} C_k^{(1)} e^{ikx} - g_1(x) \right| dx \leq \frac{\epsilon}{8} + \left[\sum_{0 \leq |k| < N_0} |C_k^{(1)}|^2 \right]^{1/2} < \frac{\epsilon}{4}. \quad (2.14)$$

Now assume that the numbers $\nu_1 < \nu_2 < \dots < \nu_{s-1}$, $N_1 < N_2 < \dots < N_{s-1}$, functions $g_1(x)$, $g_2(x), \dots, g_{s-1}(x)$, and the sets E_1, E_2, \dots, E_{s-1} are defined. We take sufficiently large natural

numbers $\nu_s > \nu_{s-1}$ and $N_s > N_{s-1}$ to satisfy

$$\frac{1}{2\pi} \left| \int_0^{2\pi} g_s(t) e^{-ikt} dt \right| < \frac{\varepsilon}{16 \cdot \sqrt{N_{s-1}}}, \quad 1 \leq s \leq q, \quad |k| < N_{s-1}, \quad (2.15)$$

$$\int_0^{2\pi} \left| \sum_{0 \leq |k| < N_s} C_k^{(s)} e^{ikx} - g_s(x) \right| dx \leq \frac{\varepsilon}{4^{s+1}}, \quad (2.16)$$

where

$$g_s(x) = \gamma_s \cdot g(\nu_s \cdot x) \cdot \chi_{\Delta_s}(x), \quad C_k^{(s)} = \frac{1}{2\pi} \int_0^{2\pi} g_s(t) e^{-ikt} dt. \quad (2.17)$$

Set

$$E_s = \{x \in \Delta_s : g_s(x) = \gamma_s\}. \quad (2.18)$$

Using the above arguments (see (2.19)–(2.21)), we conclude that the function $g_s(x)$ and the set E_s satisfy the conditions

$$|E_s| > 2\pi \cdot (1 - \varepsilon) \cdot |\Delta_s|; g_s(x) = 0, \quad x \notin \Delta_s, \quad (2.19)$$

$$\int_0^{2\pi} g_s^2(x) dx < \frac{2}{\varepsilon} \cdot |\gamma_s|^2 \cdot |\Delta_s|, \quad (2.20)$$

$$\int_0^{2\pi} \left| \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} - g_1(x) \right| dx < \frac{\varepsilon}{2^{s+1}}. \quad (2.21)$$

Thus, by induction, we can define natural numbers $\nu_1 < \nu_2 < \dots < \nu_q$, $N_1 < N_2 < \dots < N_q$, functions $g_1(x), g_2(x), \dots, g_q(x)$, and sets E_1, E_2, \dots, E_q such that conditions (2.17)–(2.19) are satisfied for all s , $1 \leq s \leq q$. We define a set E and a polynomial $P(x)$ as follows:

$$E = \bigcup_{s=1}^q E_s, \quad (2.22)$$

$$P(x) = \sum_{N_0 \leq |k| < N} C_k e^{ikx} = \sum_{s=1}^q \left[\sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right], \quad C_{-k} = \overline{C_k}, \quad (2.23)$$

where

$$C_k = C_k^{(s)} \quad \text{for } N_{s-1} \leq |k| < N_s, \quad s = 1, 2, \dots, q, \quad N = N_q - 1. \quad (2.24)$$

By Bessel's inequality and (2.5), (2.17) for all $s \in [1, q]$ we get

$$\begin{aligned} \left[\sum_{N_{s-1} \leq |k| < N_s} |C_k^{(s)}|^2 \right]^{1/2} &\leq \left[\int_0^{2\pi} g_s^2(x) dx \right]^{1/2} \\ &\leq \frac{2}{\sqrt{\varepsilon}} \cdot |\gamma_s| \cdot \sqrt{|\Delta_s|}, \quad s = 1, 2, \dots, q. \end{aligned} \quad (2.25)$$

6 Universal series by trigonometric system

From (2.5), (2.15), and (2.16), it follows that

$$|E| > 2\pi - \varepsilon. \quad (2.26)$$

Taking relations (2.1), (2.5), (2.13), (2.15), (2.21)–(2.24), we obtain

$$\int_E |P(x) - f(x)| dx \leq \sum_{s=1}^q \left[\int_E \left| \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} - g_s(x) \right| dx \right] < \varepsilon. \quad (2.27)$$

By (2.1), (2.2), (2.23)–(2.24) for any $k \in [N_0, N]$, we have

$$\begin{aligned} \sum_{N_0 \leq |k| < N} |C_k|^{2+\varepsilon} &\leq \max_{N_0 \leq k \leq N} |C_k|^\varepsilon \cdot \sum_{k=N_0}^N |C_k|^2 \\ &\leq \max_{1 \leq s \leq q} \left[\sqrt{\frac{8}{\varepsilon}} \cdot |\gamma_s| \cdot \sqrt{|\Delta_s|} \right] \cdot \sum_{s=1}^q \left[\sum_{N_{s-1} \leq |k| < N_s} |C_k^{(s)}|^2 \right] \\ &\leq \max_{1 \leq s \leq q} \left[\sqrt{\frac{8}{\varepsilon}} \cdot |\gamma_s| \cdot \sqrt{|\Delta_s|} \right] \cdot \frac{8}{\varepsilon} \cdot \sum_{s=1}^q |\gamma_s|^2 \cdot |\Delta_s| \\ &\leq \max_{1 \leq s \leq q} \left[\sqrt{\frac{8}{\varepsilon}} \cdot |\gamma_s| \cdot \sqrt{|\Delta_s|} \right] \cdot \frac{8}{\varepsilon} \cdot \left[\int_0^1 f^2(x) dx \right] < \varepsilon; \end{aligned} \quad (2.28)$$

that is, the statements (2.4a)–(2.4c) of Lemma 2.1 are satisfied. Now we will check the fulfillment of statement (2.4d) of Lemma 2.1. Let $N_0 \leq m < N$, then for some $s_0, 1 \leq s_0 \leq q$, ($N_{s_0} \leq m < N_{s_0+1}$) we will have (see (2.23) and (2.24))

$$\sum_{N_0 \leq |k| \leq m} C_k e^{ikx} = \sum_{s=1}^{s_0} \left[\sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right] + \sum_{N_{s_0-1} \leq |k| \leq m} C_k^{(s_0+1)} e^{ikx}. \quad (2.29)$$

Hence and from (2.1), (2.2), (2.5), (2.21), (2.22), and (2.25) for any measurable set $e \subset E$, we obtain

$$\begin{aligned} &\int_e \left| \sum_{N_{s-1} \leq |k| \leq m} C_k e^{ikx} \right| dx \\ &\leq \sum_{s=1}^{s_0} \left[\int_e \left| \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} - g_s(x) \right| dx \right] \\ &\quad + \sum_{s=1}^{s_0} \int_e |g_s(x)| dx + \int_e \left| \sum_{N_{s_0-1} \leq |k| \leq m} C_k^{(s_0+1)} e^{ikx} \right| dx \\ &< \sum_{s=1}^{s_0} \frac{\varepsilon}{2^{s+1}} + \int_e |f(x)| dx + \frac{2}{\sqrt{\varepsilon}} \cdot |\gamma_{s_0+1}| \cdot \sqrt{|\Delta_{s_0+1}|} < \int_e |f(x)| dx + \varepsilon. \end{aligned} \quad (2.30)$$

□

3. Proof of theorems

Proof of Theorem 1.5. Let

$$f_1(x), f_2(x), \dots, f_n(x), \quad x \in [0, 2\pi], \quad (3.1)$$

be a sequence of all step functions, values, and constancy interval endpoints of which are rational numbers. Applying lemma consecutively, we can find a sequence $\{E_s\}_{s=1}^{\infty}$ of sets and a sequence of polynomials

$$P_s(x) = \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx}, \quad (3.2)$$

$$1 = N_0 < N_1 < \dots < N_s < \dots, \quad s = 1, 2, \dots,$$

which satisfy the conditions

$$|E_s| > 1 - 2^{-2(s+1)}, \quad E_s \subset [0, 2\pi], \quad (3.3)$$

$$\int_{E_s} |P_s(x) - f_s(x)| dx < 2^{-2(s+1)}, \quad (3.4)$$

$$\sum_{N_{s-1} \leq |k| < N_s} |C_k^{(s)}|^{2+2^{-2s}} < 2^{-2s}, \quad C_{-k}^{(s)} = \overline{C_k^{(s)}}, \quad (3.5)$$

$$\max_{N_{s-1} \leq p < N_s} \left[\int_e \left| \sum_{N_{s-1} \leq |k| \leq p} C_k e^{ikx} \right| dx \right] < 2^{-2(s+1)} + \int_e |f_s(x)| dx \quad (3.6)$$

for every measurable subset e of E_s .

Denote

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} = \sum_{s=1}^{\infty} \left[\sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right], \quad (3.7)$$

where $C_k = C_k^{(s)}$ for $N_{s-1} \leq |k| < N_s$, $s = 1, 2, \dots$

Let ε be an arbitrary positive number. Setting

$$\Omega_n = \bigcap_{s=n}^{\infty} E_s, \quad n = 1, 2, \dots, \quad (3.8)$$

$$E = \Omega_{n_0} = \bigcap_{s=n_0}^{\infty} E_s, \quad n_0 = [\log_{1/2} \varepsilon] + 1, \quad (3.9)$$

$$B = \bigcup_{n=n_0}^{\infty} \Omega_n = \Omega_{n_0} \cup \left(\bigcup_{n=n_0+1}^{\infty} \Omega_n \setminus \Omega_{n-1} \right). \quad (3.10)$$

It is clear (see (3.3)) that $|B| = 2\pi$ and $|E| > 2\pi - \varepsilon$.

8 Universal series by trigonometric system

We define a function $\mu(x)$ in the following way:

$$\begin{aligned}\mu(x) &= 1 \quad \text{for } x \in E \cup ([0, 2\pi] \setminus B), \\ \mu(x) &= \mu_n \quad \text{for } x \in \Omega_n \setminus \Omega_{n-1}, \quad n \geq n_0 + 1,\end{aligned}\tag{3.11}$$

where

$$\mu_n = \left[2^{4n} \cdot \prod_{s=1}^n h_s \right]^{-1},\tag{3.12}$$

$$h_s = \|f_s(x)\|_C + \max_{N_{s-1} \leq p < N_s} \left\| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right\|_C + 1,\tag{3.13}$$

where

$$\|g(x)\|_C = \max_{x \in [0, 2\pi]} |g(x)|,\tag{3.14}$$

$g(x)$ is a continuous function on $[0, 2\pi]$.

From (3.5), (3.7)–(3.12), we obtain the following.

(A) $0 < \mu(x) \leq 1$, $\mu(x)$ is a measurable function and

$$|\{x \in [0, 2\pi] : \mu(x) \neq 1\}| < \varepsilon.\tag{3.15}$$

(B) $\sum_{k=1}^{\infty} |C_k|^q < \infty$ for all $q > 2$.

Hence, obviously, we have

$$\lim_{k \rightarrow \infty} C_k = 0.\tag{3.16}$$

It follows from (3.9)–(3.12) that for all $s \geq n_0$ and $p \in [N_{s-1}, N_s)$,

$$\begin{aligned}\int_{[0, 2\pi] \setminus \Omega_s} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu(x) dx &= \sum_{n=s+1}^{\infty} \left[\int_{\Omega_n \setminus \Omega_{n-1}} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu_n dx \right] \\ &\leq \sum_{n=s+1}^{\infty} 2^{-4n} \left[\int_0^{2\pi} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| h_s^{-1} dx \right] < 2^{-4s}.\end{aligned}\tag{3.17}$$

By (3.4), (3.9)–(3.12) for all $s \geq n_0$, we have

$$\begin{aligned}&\int_0^{2\pi} |P_s(x) - f_s(x)| \mu(x) dx \\ &= \int_{\Omega_s} |P_s(x) - f_s(x)| \mu(x) dx + \int_{[0, 2\pi] \setminus \Omega_s} |P_s(x) - f_s(x)| \mu(x) dx \\ &= 2^{-2(s+1)} + \sum_{n=s+1}^{\infty} \left[\int_{\Omega_n \setminus \Omega_{n-1}} |P_s(x) - f_s(x)| \mu_n dx \right]\end{aligned}$$

$$\begin{aligned}
&\leq 2^{-2(s+1)} + \sum_{n=s+1}^{\infty} 2^{-4s} \left[\int_0^{2\pi} \left(|f_s(x)| + \left| \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right| \right) h_s^{-1} dx \right] \\
&< 2^{-2(s+1)} + 2^{-4s} < 2^{-2s}.
\end{aligned} \tag{3.18}$$

Taking relations (3.6), (3.9)–(3.12), and (3.17) into account we obtain that for all $p \in [N_{s-1}, N_s]$ and $s \geq n_0 + 1$,

$$\begin{aligned}
&\int_0^{2\pi} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu(x) dx \\
&= \int_{\Omega_s} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu(x) dx + \int_{[0, 2\pi] \setminus \Omega_s} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu(x) dx \\
&< \sum_{n=n_0+1}^s \left[\int_{\Omega_n \setminus \Omega_{n-1}} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| dx \right] \cdot \mu_n + 2^{-4s} \\
&< \sum_{n=n_0+1}^s \left(2^{-2(s+1)} + \int_{\Omega_n \setminus \Omega_{n-1}} |f_s(x)| dx \right) \mu_n + 2^{-4s} \\
&= 2^{-2(s+1)} \cdot \sum_{n=n_0+1}^s \mu_n + \int_{\Omega_s} |f_s(x)| \mu(x) dx + 2^{-4s} \\
&< \int_0^{2\pi} |f_s(x)| \mu(x) dx + 2^{-4s}.
\end{aligned} \tag{3.19}$$

Let $f(x) \in L_{\mu}^1[0, 2\pi]$, that is, $\int_0^{2\pi} |f(x)| \mu(x) dx < \infty$.

It is easy to see that we can choose a function $f_{\nu_1}(x)$ from the sequence (3.1) such that

$$\int_0^{2\pi} |f(x) - f_{\nu_1}(x)| \mu(x) dx < 2^{-2}, \quad \nu_1 > n_0 + 1. \tag{3.20}$$

Hence, we have

$$\int_0^{2\pi} |f_{\nu_1}(x)| \mu(x) dx < 2^{-2} + \int_0^{2\pi} |f(x)| \mu(x) dx. \tag{3.21}$$

From (2.1), (A), (3.18), and (3.20), we obtain with $m_1 = 1$,

$$\begin{aligned}
&\int_0^{2\pi} |f(x) - [P_{\nu_1}(x) + C_{m_1} e^{im_1 x}]| \mu(x) dx \\
&\leq \int_0^{2\pi} |f(x) - f_{\nu_1}(x)| \mu(x) dx + \int_0^{2\pi} |f_{\nu_1}(x) - P_{\nu_1}(x)| \mu(x) dx \\
&\quad + \int_0^{2\pi} |C_{m_1} e^{im_1 x}| \mu(x) dx < 2 \cdot 2^{-2} + 2\pi \cdot |C_{m_1}|.
\end{aligned} \tag{3.22}$$

10 Universal series by trigonometric system

Assume that numbers $\nu_1 < \nu_2 < \dots < \nu_{q-1}$, $m_1 < m_2 < \dots < m_{q-1}$ are chosen in such a way that the following condition is satisfied:

$$\int_0^{2\pi} \left| f(x) - \sum_{s=1}^j [P_{\nu_s}(x) + C_{m_s} e^{im_s x}] \right| \mu(x) dx < 2 \cdot 2^{-2j} + 2\pi \cdot |C_{m_j}|, \quad 1 \leq j \leq q-1. \quad (3.23)$$

We choose a function $f_{\nu_q}(x)$ from the sequence (3.1) such that

$$\int_0^{2\pi} \left| \left(f(x) - \sum_{s=1}^{q-1} [P_{\nu_s}(x) + C_{m_s} e^{im_s x}] \right) - f_{\nu_q}(x) \right| \mu(x) dx < 2^{-2q}, \quad (3.24)$$

where $\nu_q > \nu_{q-1}$; $\nu_q > m_{q-1}$

This, with (3.23), implies

$$\int_0^{2\pi} |f_{\nu_q}(x)| \mu(x) dx < 2^{-2q} + 2 \cdot 2^{-2(q-1)} + 2\pi \cdot |C_{m_{q-1}}| = 9 \cdot 2^{-2q} + 2\pi \cdot |C_{m_{q-1}}|. \quad (3.25)$$

By (3.18), (3.19), and (3.25) we obtain

$$\int_0^{2\pi} |f_{\nu_q}(x) - P_{\nu_q}(x)| \mu(x) dx < 2^{-2\nu_q}, \quad (3.26)$$

$$P_{\nu_q}(x) = \sum_{N_{\nu_q-1} \leq |k| < N_{\nu_q}} C_k^{(\nu_q)} e^{ikx},$$

$$\max_{N_{\nu_q-1} \leq p < N_{\nu_q}} \int_0^{2\pi} \left| \sum_{k=N_{\nu_q-1}}^p C_k^{(\nu_q)} e^{ikx} \right| \mu(x) dx < 10 \cdot 2^{-2q} + 2\pi \cdot |C_{m_{q-1}}|. \quad (3.27)$$

Denote

$$m_q = \min \left\{ n \in N : n \notin \left\{ \left\{ \{k\}_{k=N_{\nu_s-1}}^{N_{\nu_s}-1} \right\}_{s=1}^q \cup \{m_s\}_{s=1}^{q-1} \right\} \right\}. \quad (3.28)$$

From (2.1), (A), (3.24), and (3.26), we have

$$\begin{aligned} & \int_0^{2\pi} \left| f(x) - \sum_{s=1}^q [P_{\nu_s}(x) + C_{m_s} e^{im_s x}] \right| \mu(x) dx \\ & \leq \int_0^{2\pi} \left| \left(f(x) - \sum_{s=1}^{q-1} [P_{\nu_s}(x) + C_{m_s} e^{im_s x}] \right) - f_{\nu_q}(x) \right| \mu(x) dx \end{aligned}$$

$$\begin{aligned}
 & + \int_0^{2\pi} |f_{\nu_q}(x) - P_{\nu_q}(x)| \mu(x) dx \\
 & + \int_0^{2\pi} |C_{m_q} e^{im_q x}| \mu(x) dx < 2 \cdot 2^{-2q} + 2\pi \cdot |C_{m_q}|.
 \end{aligned} \tag{3.29}$$

Thus, by induction we, on q , can choose from series (3.7) a sequence of members

$$C_{m_q} e^{im_q x}, \quad q = 1, 2, \dots, \tag{3.30}$$

and a sequence of polynomials

$$P_{\nu_q}(x) = \sum_{N_{\nu_{q-1}} \leq |k| < N_{\nu_q}} C_k^{(\nu_q)} e^{ikx}, \quad N_{n_q-1} > N_{n_{q-1}}, \quad q = 1, 2, \dots \tag{3.31}$$

such that conditions (3.27)–(3.29) are satisfied for all $q \geq 1$.

Taking account the choice of $P_{\nu_q}(x)$ and $C_{m_q} e^{im_q x}$ (see (3.28) and (3.31)), we conclude that the series

$$\sum_{q=1}^{\infty} \left[\sum_{N_{\nu_{q-1}} \leq |k| < N_{\nu_q}} C_k^{(\nu_q)} e^{ikx} + C_{m_q} e^{iqx} \right] \tag{3.32}$$

is obtained from the series (3.7) by rearrangement of members. Denote this series by $\sum C_{\sigma(k)} e^{i\sigma(k)x}$.

It follows from (3.16), (3.27), and (3.29) that the series $\sum C_{\sigma(k)} e^{i\sigma(k)x}$ converges to the function $f(x)$ in the metric $L^1_{\mu}[0, 2\pi]$, that is, the series (3.7) is universal with respect to rearrangements (see Definition 1.1). \square

Proof of Theorem 1.5. Applying Lemma 2.1 consecutively, we can find a sequence $\{E_s\}_{s=1}^{\infty}$ of sets and a sequence of polynomials

$$P_s(x) = \sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx}, \quad C_{-k}^{(s)} = \overline{C_k^{(s)}}, \tag{3.33}$$

$$1 = N_0 < N_1 < \dots < N_s < \dots, \quad s = 1, 2, \dots,$$

which satisfy the conditions

$$|E_s| > 1 - 2^{-2(s+1)}, \quad E_s \subset [0, 2\pi], \tag{3.34}$$

$$\sum_{N_{s-1} \leq |k| < N_s} |C_k^{(s)}|^{2+2^{-2s}} < 2^{-2s}, \tag{3.35}$$

$$\int_{E_n} \left| f_n(x) - \sum_{s=1}^n P_s(x) \right| dx < 2^{-n}, \quad n = 1, 2, \dots, \tag{3.36}$$

where $\{f_n(x)\}_{n=1}^{\infty}$, $x \in [0, 2\pi]$, is a sequence of all step functions, values, and constancy interval endpoints of which are rational numbers.

12 Universal series by trigonometric system

Denote

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} = \sum_{s=1}^{\infty} \left[\sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right], \quad (3.37)$$

where $C_k = C_k^{(s)}$ for $N_{s-1} \leq |k| < N_s$, $s = 1, 2, \dots$.

It is clear (see (3.35)) that

$$\sum_{k=1}^{\infty} |C_k|^q < \infty \quad \forall q > 2. \quad (3.38)$$

Repeating reasoning of Theorem 1.4, a weighted function $\mu(x)$, $0 < \mu(x) \leq 1$, can be constructed so that the following condition is satisfied:

$$\int_0^{2\pi} \left| f_n(x) - \sum_{s=1}^n P_s(x) \right| \cdot \mu(x) dx < 2^{-2n}, \quad n = 1, 2, \dots \quad (3.39)$$

For any function $f(x) \in L_{\mu}^1[0, 1]$, we can choose a subsystem $\{f_{n_\nu}(x)\}_{\nu=1}^{\infty}$ from the sequence (3.1) such that

$$\int_0^{2\pi} |f(x) - f_{n_\nu}(x)| \mu(x) dx < 2^{-2\nu}. \quad (3.40)$$

From (3.37)–(3.40), we conclude

$$\begin{aligned} \int_0^{2\pi} \left| f(x) - \sum_{|k| \leq M_\nu} C_k e^{ikx} \right| \mu(x) dx &= \int_0^{2\pi} \left| f(x) - \sum_{s=1}^{n_\nu} \left[\sum_{N_{s-1} \leq |k| < N_s} C_k^{(s)} e^{ikx} \right] \right| \mu(x) dx \\ &\leq \int_0^{2\pi} |f(x) - f_{n_\nu}(x)| \cdot \mu(x) dx \\ &\quad + \int_0^{2\pi} \left| f_{n_\nu}(x) - \sum_{s=1}^{n_\nu} P_s(x) \right| \cdot \mu(x) dx < 2^{-2k} + 2^{-2\nu k}, \end{aligned} \quad (3.41)$$

where $M_\nu = N_{n_\nu} - 1$.

Thus, the series (3.37) is universal in $L_{\mu}^1[0, 1]$ in the sense of usual (see Definition 1.2).

Theorem 1.5 is proved. \square

Acknowledgments

The author was supported in part by Grant 01-000 from the Government of Armenia. The author thanks Professor M. G. Grigorian for his attention to this paper.

References

- [1] F. G. Arutjunjan, *Representation of functions by multiple series*, Akademiya Nauk Armyanskoï SSR. Doklady **64** (1977), no. 2, 72–76.
- [2] S. A. Episkoposian, *On the existence of universal series by trigonometric system*, Journal of Functional Analysis **230** (2006), no. 1, 169–183.
- [3] M. G. Grigorian, *On the representation of functions by orthogonal series in weighted L^p spaces*, Studia Mathematica **134** (1999), no. 3, 207–216.
- [4] M. G. Grigorian and S. A. Episkoposian, *Representation of functions in the weighted space $L^1_\mu[0, 2\pi]$ by trigonometric and Walsh series*, Analysis Mathematica **27** (2001), no. 4, 261–277.
- [5] V. Ya. Kozlov, *On complete systems of orthogonal functions*, Matematicheskii Sbornik **26(68)** (1950), 351–364.
- [6] D. E. Menshov, *On the partial sums of trigonometric series*, Matematicheskii Sbornik **20** (1947), 197–238.
- [7] W. Orlicz, *Über die unabhängig von der Anordnung fast überall kniwergernten Reihen*, Bulletin de l'Académie Polonaise des Science **81** (1927), 117–125.
- [8] A. A. Talaljan, *On the convergence almost everywhere of subsequences of partial sums of general orthogonal series*, Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya **10** (1957), no. 3, 17–34.
- [9] ———, *Series universal with respect to rearrangement*, Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya **24** (1960), 567–604.
- [10] P. L. Ul'yanov, *Unconditional convergence and summability*, Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya **22** (1958), 811–840.

S. A. Episkoposian: Chair of Higher Mathematics, Faculty of Physics, Yerevan State University,
 Alex Manoogian street 1, Yerevan 375049, Armenia
 E-mail address: sergoep@ysu.am



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

