

Corrigendum

Corrigendum to “Effects of Number of Repetitions and Number of Hours of Shaping Practice during Constraint-Induced Movement Therapy: A Randomized Controlled Trial”

Auwal Abdullahi 

Department of Physiotherapy, Bayero University Kano, Kano, Kano State, Nigeria

Correspondence should be addressed to Auwal Abdullahi; aabdullahi.pth@buk.edu.ng

Received 28 July 2020; Accepted 28 July 2020; Published 24 August 2020

Copyright © 2020 Auwal Abdullahi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Effects of Number of Repetitions and Number of Hours of Shaping Practice during Constraint-Induced Movement Therapy: A Randomized Controlled Trial” [1], there was an error in Table 1 in which the

incorrect data were given for sex, type of stroke, hand dominance, and side affected. The author apologises for this error and the corrected table is shown below.

TABLE 1: Baseline characteristics of the study participants.

Variable	Control ($n = 12$)	Group			Statistics	p
		Modified CIMT ($n = 13$)	300 Rep ($n = 12$)	600 Rep ($n = 11$)		
Age	58.83 ± 10.57	54.62 ± 6.00	59.42 ± 13.93	57.60 ± 10.27		
Time since stroke	19.89 ± 7.20	14.75 ± 4.46	21.67 ± 6.38	13.50 ± 7.39		
Sex (M/F)	9/3	9/4	4/8	5/6		
Type of stroke (I/H)	8/4	10/3	11/1	7/4		
Hand dominance (R/L)	12/0	12/1	11/1	10/1		
Side affected (R/L)	7/5	9/4	7/5	8/3		
MAL (how well)	2.13 ± 1.32	1.62 ± 0.73	2.30 ± 1.16	1.51 ± 0.94	$F = 1.577$	0.208
MAL (amount of use)	1.91 ± 1.37	1.75 ± 0.54	2.31 ± 1.19	1.18 ± 0.77	$F = 2.433$	0.078
WMFT	2.22 ± 1.19	2.17 ± 0.82	1.05 ± 0.30	1.99 ± 0.52	$F = 0.524$	0.668
FM	31.42 ± 16.21	34.00 ± 14.15	35.17 ± 13.02	30.45 ± 10.76	$F = 0.299$	0.826
UPSET	4.75 ± 3.26	4.58 ± 2.21	4.74 ± 1.88	4.56 ± 1.81	$F = 0.021$	0.996

R/L = right/left; I/H = ischaemic/haemorrhagic; M/F = male/female.

References

- [1] A. Abdullahi, "Effects of number of repetitions and number of hours of shaping practice during constraint-induced movement therapy: a randomized controlled trial," *Neurology Research International*, vol. 2018, Article ID 5496408, 9 pages, 2018.